Gramin Krishi Mausam Sewa

Agromet Advisory Bulletin for SOUTH-ANDAMANS District

Central Agricultural Research Institute

Port Blair, Andaman & Nicobar Islands

Agromet Advisory Bulletin

Date : 2026-11-03

Weather Forecast of District SOUTH-ANDAMANS (West Bengal) Issued On : 2026-11-03 (Valid Till 08:30 IST of the next 5 days)

Parameter	2026-11-04	2026-11-05	2026-11-06	2026-11-07	2026-11-08
Rainfall(mm)	3.0	5.0	7.0	12.0	8.0
Tmax(°C)	30.0	30.0	30.0	30.0	30.0
Tmin(°C)	25.0	25.0	25.0	25.0	25.0
RH-I(%)	89	88	87	87	81
RH-II(%)	73	72	70	72	61
Wind Speed(kmph)	11	10	5	5	5
Wind Direction(Degree)	32	49	48	23	34
Cloud Cover(Octa)	5	5	6	3	8
Warning	No Warning	No Warning	No Warning	No Warning	No Warning

Forecast Summary:

The weather forecast for the next five days, starting from 04 November 2026, indicates that light rainfall is expected, with rainfall amounts ranging from 3.00 to 12.00 mm and cumulative rainfall expected to be 35.00 mm. The maximum temperature is expected to be around 30.0°C, while the minimum temperature will be around 25.0°C. Maximum relative humidity is likely to range from 81% to 89%, and minimum relative humidity from 61% to 73%. Wind speed is expected to vary between 5 and 11 kmph, with wind directions varying from north-northeast (23°) to northeast (49°). Cloud conditions are expected to remain partly cloudy to cloudy.

Weather Warnings (Valid Till 08:30 IST of the next day)

There are no weather warnings for upcoming five days.

Likely impacts of weather warnings on Agriculture and associated Agromet advisories

Since no warnings have been issued, no significant damage is expected.

General Advisory:

Considering the forecast of light rainfall, farmers are advised to use the expected rainfall for planning irrigation and avoid irrigation where adequate soil moisture is available. Fertilizer and plant protection applications may be taken up during suitable rain-free periods after checking the latest rainfall forecast. Monitor crops regularly for pests and diseases, particularly under humid conditions. Keep livestock in clean, dry and well-ventilated shelters and protect feed and fodder from rain and moisture. Conserve rainwater where possible and maintain farm ponds and other water storage structures. Follow weather updates through the Meghdoot app and lightning alerts through the Damini app.

SMS Advisory:

To prevent African swine fever, restrict outsiders' entry into pig farms. Keep the farm, pens and equipment clean and disinfect regularly. Provide clean water and balanced feed.

Crop Specific Advisory:

Crop (Stage)	Crop Specific Advisory
RICE (Physiological Maturity)	Considering the forecast of light rainfall during the coming days, farmers are advised to prepare for harvesting rice crops that have attained physiological maturity and undertake harvesting during clear and sunny weather conditions. Drain excess water from rice fields approaching maturity to prevent water stagnation and facilitate proper crop maturation and timely harvesting.Keep the harvested produce in a safe and dry place to protect it from moisture. Always check the latest weather forecast before undertaking harvesting operations.
NUTMEG (Flowering)	Considering the emergence of new flushes in clove and nutmeg, farmers are advised to clean the basins of the plants and apply coconut husk mulch to conserve soil moisture and support bud development.

Horticulture Specific Advisory:

Horticulture (Stage)	Horticulture Specific Advisory
BLACK PEPPER (Vegetative)	In black pepper, if yellowing of leaf veins is observed, farmers are advised to apply neem cake @ 500 g per plant and drench the soil with 1% Bordeaux mixture. For preparing 100 litres of 1% Bordeaux mixture, dissolve 1 kg copper sulphate in 50 litres of water and 1 kg fresh lime separately in 50 litres of water; slowly mix the copper sulphate solution into the lime suspension with continuous stirring, filter and use the mixture fresh.
BRINJAL (Fruit formation)	Farmers are advised to regularly inspect brinjal fields. If Bacterial wilt symptoms observed, remove and destroy the infected plants to prevent the spread of the disease. Practise Crop rotaion with non solanaceous crops Grow bacterial wilt-resistant brinjal varieties such as Dweep CARI Brinjal-1. Apply 5% bleaching powder as a soil drench and use streptocycline 2% for seed treatment.
COCONUT	Farmers who have not yet applied post-monsoon fertilizers are advised to apply well-decomposed organic manure along with 50% of the recommended dose of fertilizers (RDF) to coconut, arecanut and other perennial crops. Apply fertilizers around the palm basin and incorporate them into the soil during rain-free periods to minimize nutrient loss. Always check the latest weather forecast before fertilizer application and avoid applying fertilizers when rainfall is expected.
GINGER (Grand growth)	As ginger, turmeric and mango ginger crops are in the rhizome development stage, farmers are advised to carry out earthing up and mulching to cover exposed rhizomes, conserve soil moisture and promote healthy rhizome development. Use organic mulch such as dry leaves or crop residues, and ensure proper drainage to prevent waterlogging.

Live Stock Specific Advisory:

Live Stock	Live Stock Specific Advisory
PIG	Considering the risk of African Swine Fever (ASF) transmission, farmers are advised to restrict the entry of outsiders into pig farms and maintain proper farm hygiene by keeping the surroundings clean and regularly cleaning and disinfecting pig pens and equipment. Provide pigs with good-quality, balanced feed and clean drinking water in adequate quantities, and avoid feeding contaminated feed or water. Healthy pigs should be dewormed at regular intervals as recommended by the local veterinarian and provided with appropriate vitamin and mineral supplements to maintain their health and improve overall productivity.
COW	Farmers are advised to provide livestock with clean and properly filtered drinking water. Animals should not be allowed to bathe or wallow in stagnant water to prevent the spread of waterborne diseases. Animal dipping should be carried out during daytime under suitable weather conditions. Regular

Live Stock	Live Stock Specific Advisory
	deworming should also be undertaken using appropriate deworming medicines as per veterinary advice.

Likely impacts of weather warnings (General)

There is no weather warning for the upcoming five days. No significant weather-related impact or damage is expected during this period.

Impact based advisories (General)

No impact-based advisory is provided as there is no weather warning.Carry out routine household, workplace and outdoor activities as planned. Keep checking the latest weather forecast and warnings and take necessary precautions if the weather situation changes.

Farmers are advised to download Unified “Mausam” and "Meghdoot" android application on mobile for Weather forecast and weather based Agromet Advisories and "Damini" android application for forecast of Thunderstorm and lightening.

Mausam MobileApp link: https://play.google.com/store/apps/details/

Meghdoot MobileApp link: https://play.google.com/store/apps/details

Damini MobileApp link : https://play.google.com/store/apps/details

ग्रामीण कृषि मौसम सेवा
भारत मौसम विज्ञान विभाग
केंद्रीय द्ववीपीय कृषि अनुसंधान संसथान
पोर्टब्लेयर, अंडमान तथा निकोबार द्ववीपसमूह

मौसम आधारित कृषि परामर्श सेवाएं

दिनांक : 03-11-2026

SOUTH-ANDAMANS(पश्चिम वङ्ग) के मौसम का पूर्वानुमान - जारी करनेका दिन :2026-11-03 (अगले 5 दिनों के 8:30 IST तक वैध)

मौसम कारक	2026-11-04	2026-11-05	2026-11-06	2026-11-07	2026-11-08
वर्षा (मिमी)	3.0	5.0	7.0	12.0	8.0
अधिकतम तापमान(से.)	30.0	30.0	30.0	30.0	30.0
न्यूनतम तापमान(से.)	25.0	25.0	25.0	25.0	25.0
अधिकतम सापेक्षिक आर्द्रता (%)	89	88	87	87	81
न्यूनतम सापेक्षिक आर्द्रता (%)	73	72	70	72	61
हवा की गति (किमी प्रति घंटा)	11	10	5	5	5
पवन दिशा (डिग्री)	32	49	48	23	34
क्लाउड कवर (ओक्टा)	5	5	6	3	8
चेतावनी	कोई चेतावनी नहीं	कोई चेतावनी नहीं	कोई चेतावनी नहीं	कोई चेतावनी नहीं	कोई चेतावनी नहीं

पूर्वानुमान सारांश:

04 नवंबर 2026 से शुरू होने वाले अगले पाँच दिनों के दौरान हल्की वर्षा होने की संभावना है। वर्षा की मात्रा 3.00 से 12.00 मिमी के बीच रहने तथा कुल संचयी वर्षा 35.00 मिमी होने की संभावना है। अधिकतम तापमान लगभग 30.0°C तथा न्यूनतम तापमान लगभग 25.0°C रहने की संभावना है। अधिकतम सापेक्ष आर्द्रता 81% से 89% तथा न्यूनतम सापेक्ष आर्द्रता 61% से 73% के बीच रहने की संभावना है। हवा की गति 5 से 11 किमी प्रति घंटा रहने तथा हवा की दिशा उत्तर-उत्तर-पूर्व (23°) से उत्तर-पूर्व (49°) के बीच परिवर्तित होने की संभावना है। आसमान में आंशिक रूप से बादल छाए रहने से लेकर सामान्यतः बादल छाए रहने की संभावना है।

मौसम चेतावनियाँ (अगले दिन के 08:30 IST तक मान्य):

आगामी पाँच दिनों के लिए कोई मौसम चेतावनी नहीं है।

मौसम चेतावनियों के कृषि पर संभावित प्रभाव और संबंधित एग्रोमेट सलाह:

किसी मौसम चेतावनी के जारी नहोनेके कारण कोई महत्वपूर्ण क्षतिहोने की संभावना नहीं है।

सामान्य सलाहकार:

हल्की वर्षा के मौसम पूर्वानुमान को ध्यान में रखते हुए किसानों को सलाह दी जाती है कि संभावित वर्षा का उपयोग सिंचाई प्रबंधन की योजना बनाने में करें तथा जहां मिट्टी में पर्याप्त नमी उपलब्ध हो, वहां सिंचाई से बचें। उर्वरकों एवं

पौध संरक्षण रसायनों का प्रयोग नवीनतम मौसम पूर्वानुमान की जानकारी लेने के बाद उपयुक्त वर्षा रहित अवधि में ही करें। आर्द्र परिस्थितियों में कीट एवं रोगों की संभावना को देखते हुए फसलों की नियमित निगरानी करें। पशुओं को स्वच्छ, सूखे एवं हवादार आश्रय स्थलों में रखें तथा चारा एवं पशु आहार को वर्षा एवं नमी से सुरक्षित रखें। वर्षा जल का अधिकतम संचयन करें तथा खेतों में बने तालाबों एवं अन्य जल भंडारण संरचनाओं का उचित रखरखाव करें। मौसम संबंधी जानकारी के लिए मेघदूत (Meghdoot) ऐप तथा आकाशीय बिजली से संबंधित चेतावनियों के लिए दामिनी (Damini) ऐप का नियमित उपयोग करें।

लघु संदेश सलाहकार:

African Swine Fever से बचाव हेतु सूअर फार्म में बाहरी व्यक्तियों का प्रवेश सीमित रखें। फार्म, बाड़ों एवं उपकरणों की नियमित सफाई एवं कीटाणुशोधन करें। सूअरों को स्वच्छ पानी एवं संतुलित आहार प्रदान करें।

फ़सल विशिष्ट सलाह:

फ़सल	फ़सल विशिष्ट सलाह
चावल	आगामी दिनों में हल्की वर्षा की संभावना को देखते हुए, किसानों को सलाह दी जाती है कि जिन धान की फसलों ने शारीरिक परिपक्वता (Physiological Maturity) प्राप्त कर ली है, उनकी कटाई की तैयारी करें तथा कटाई का कार्य साफ एवं धूप वाले मौसम में करें। परिपक्वता के निकट पहुँच चुकी धान की फसलों वाले खेतों से अतिरिक्त पानी निकाल दें, ताकि खेत में जलभराव न हो तथा फसल के उचित परिपक्वन और समय पर कटाई में सुविधा हो। कटी हुई उपज को नमी से बचाने के लिए सुरक्षित एवं सूखे स्थान पर रखें। कटाई का कार्य करने से पहले हमेशा नवीनतम मौसम पूर्वानुमान की जानकारी अवश्य प्राप्त करें।
जायफल	लौंग एवं जायफल के पौधों में नई कोंपलों के निकलने को ध्यान में रखते हुए किसानों को सलाह दी जाती है कि पौधों के थालों की सफाई करें तथा मिट्टी में नमी बनाए रखने और कलियों के समुचित विकास को बढ़ावा देने के लिए नारियल की भूसी से मल्चिंग करें।

बागवानी विशिष्ट सलाह:

बागवानी	बागवानी विशिष्ट सलाह
काली मिर्च	काली मिर्च में पत्तियों की शिराओं के पीले पड़ने के लक्षण दिखाई देने पर किसानों को सलाह दी जाती है कि प्रत्येक पौधे में 500 ग्राम नीम की खली डालें तथा 1% बोर्डो मिश्रण से मिट्टी की सिंचाई (ड्रेंचिंग) करें। 100 लीटर 1% बोर्डो मिश्रण तैयार करने के लिए 1 किलोग्राम कॉपर सल्फेट को 50 लीटर पानी में तथा 1 किलोग्राम ताजे बुझा हुए चूने को अलग से 50 लीटर पानी में घोलें। इसके बाद लगातार हिलाते हुए कॉपर सल्फेट के घोल को धीरे-धीरे चूने के घोल में मिलाएं। मिश्रण को छानकर ताजा ही उपयोग करें।
बैंगन	किसानों को सलाह दी जाती है कि बैंगन के खेतों का नियमित निरीक्षण करें। यदि Bacterial wilt के लक्षण दिखाई दें, तो रोग के प्रसार को रोकने के लिए संक्रमित पौधों को उखाड़कर नष्ट कर दें। अन्य कुलों की सब्जियों वाली फसलों के साथ फसल चक्र अपनाएँ तथा जीवाणु म्लानि प्रतिरोधी बैंगन की किस्मों, जैसे Dweep CARI Brinjal-1, की खेती करें। मिट्टी में 5% ब्लीचिंग पाउडर (Bleaching Powder) का घोल डालें तथा बीज उपचार के लिए 2% Streptocyclin का उपयोग करें।
नारियल	जिन किसानों ने अभी तक मानसूनोत्तर उर्वरकों का प्रयोग नहीं किया है, उन्हें सलाह दी जाती है कि वे नारियल, सुपारी एवं अन्य बहुवर्षीय फसलों में अच्छी तरह सड़ी हुई जैविक खाद के साथ अनुशंसित उर्वरक मात्रा (RDF) का 50% प्रयोग करें। उर्वरकों को पौधों के थालों के चारों ओर डालकर मिट्टी में अच्छी तरह मिला दें। पोषक तत्वों के नुकसान को कम करने के लिए उर्वरकों का प्रयोग वर्षा रहित अवधि में ही करें। उर्वरकों के प्रयोग से पूर्व नवीनतम मौसम पूर्वानुमान अवश्य देखें तथा वर्षा की संभावना होने पर उर्वरकों का प्रयोग न करें।
अदरक	अदरक, हल्दी एवं आम-अदरक की फसलें वर्तमान में प्रकंद विकास अवस्था में हैं। अतः किसानों को सलाह दी जाती है कि खुले हुए प्रकंदों को मिट्टी से ढकने, मृदा में नमी बनाए रखने तथा प्रकंदों के समुचित विकास को बढ़ावा देने के लिए मिट्टी चढ़ाने (अर्थिंग-अप) एवं पलवार (मल्चिंग) की प्रक्रिया अपनाएं। इसके लिए सूखी पत्तियों या फसल अवशेषों जैसे जैविक पदार्थों का उपयोग करें तथा खेत में जलभराव रोकने के लिए उचित जल निकासी की व्यवस्था सुनिश्चित करें।

पशुपालन विशिष्ट सलाह:

पशुपालन	पशुपालन विशिष्ट सलाह
सूअर	अफ्रीकन स्वाइन फीवर (ASF) के संक्रमण के जोखिम को देखते हुए, किसानों को सलाह दी जाती है कि वे सूअरों के फार्म में बाहरी व्यक्तियों के प्रवेश को प्रतिबंधित करें तथा फार्म परिसर को स्वच्छ रखें। सूअरशालाओं एवं उपयोग में आने वाले उपकरणों की नियमित रूप से सफाई एवं कीटाणुशोधन करें। सूअरों को पर्याप्त मात्रा में अच्छी गुणवत्ता वाला संतुलित आहार एवं स्वच्छ पेयजल उपलब्ध कराएं तथा दूषित आहार या पानी देने से बचें। स्वस्थ सूअरों को स्थानीय पशु चिकित्सक की सलाह के अनुसार नियमित अंतराल पर कृमिनाशक दवा दें तथा उनके स्वास्थ्य और उत्पादकता को बनाए रखने के लिए उचित मात्रा में विटामिन एवं खनिज पूरक प्रदान करें।
गाय	पशुपालकों को सलाह दी जाती है कि पशुओं को स्वच्छ एवं उचित रूप से छना हुआ पेयजल उपलब्ध कराएं। जलजनित रोगों के प्रसार को रोकने के लिए पशुओं को ठहरे हुए पानी में न नहाने दें और न ही उन्हें उसमें लोटने दें। पशुओं का डिपिंग (परजीवी नियंत्रण हेतु औषधीय घोल में स्नान) दिन के समय अनुकूल मौसम परिस्थितियों में करें। पशुओं को पशु चिकित्सक की सलाह के अनुसार उपयुक्त कृमिनाशक दवाओं से नियमित रूप से कृमिनाशन कराएं।

मौसम चेतावनियों के संभावित प्रभाव (सामान्य):

आगामी पाँच दिनों के लिए कोई मौसम चेतावनी नहीं है। इस अवधिके दौरान मौसम से संबंधित किसी महत्वपूर्ण प्रभाव याक्षतिकी संभावना नहीं है।

प्रभाव आधारित सलाह (सामान्य):

मौसम संबंधी कोई चेतावनी नहीं होने के कारण कोई प्रभाव-आधारित परामर्श नहीं है।घर, कार्यस्थलऔर बाहर के नियमित कार्य योजनानु सार करें।मौसम की स्थिति में बदलाव होने परआवश्यक सावधानियाँ बरतनेके लिए नवीनतम मौसम पूर्वानुमानऔर चेतावनियों की जानकारी लेते रहें।

Farmers are advised to download Unified ‘Mausam’ and "Meghdoot" android application on mobile for Weather forecast and weather based Agromet Advisories and "Damini" android application for forecast of Thunderstorm and lightening.

Mausam MobileApp link: https://play.google.com/store/apps/details/

Meghdoot MobileApp link: https://play.google.com/store/apps/details

Damini MobileApp link : https://play.google.com/store/apps/details

Gramin Krishi Mausam Sewa

Agromet Advisory Bulletin for N-AND-M-ANDAMAN District

Central Agricultural Research Institute

Port Blair, Andaman & Nicobar Islands

Agromet Advisory Bulletin

Date : 2026-11-03

Weather Forecast of District N-AND-M-ANDAMAN (West Bengal) Issued On : 2026-11-03 (Valid Till 08:30 IST of the next 5 days)

Parameter	2026-11-04	2026-11-05	2026-11-06	2026-11-07	2026-11-08
Rainfall(mm)	3.0	3.0	7.0	8.0	5.0
Tmax(°C)	30.0	30.0	30.0	30.0	30.0
Tmin(°C)	25.0	25.0	25.0	25.0	25.0
RH-I(%)	89	89	88	86	79
RH-II(%)	69	71	68	70	57
Wind Speed(kmph)	10	7	3	5	6
Wind Direction(Degree)	9	32	41	13	32
Cloud Cover(Octa)	5	3	4	3	6
Warning	No Warning	No Warning	No Warning	No Warning	No Warning

Forecast Summary:

The weather forecast for the next five days, starting from 04 November 2026, indicates that light rainfall is expected, with rainfall amounts ranging from 3.00 to 8.00 mm and cumulative rainfall expected to be 26.00 mm. The maximum temperature is expected to be around 30.0°C, while the minimum temperature will be around 25.0°C. Maximum relative humidity is likely to range from 79% to 89%, and minimum relative humidity from 57% to 71%. Wind speed is expected to vary between 3 and 10 kmph, with wind directions varying from north (9°) to northeast (41°). Cloud conditions are expected to remain partly cloudy to generally cloudy.

Weather Warnings (Valid Till 08:30 IST of the next day)

There are no weather warnings for upcoming five days.

Likely impacts of weather warnings on Agriculture and associated Agromet advisories

Since no warnings have been issued, no significant damage is expected.

General Advisory:

Considering the forecast of light rainfall, farmers are advised to use the expected rainfall for planning irrigation and avoid irrigation where adequate soil moisture is available. Fertilizer and plant protection applications may be taken up during suitable rain-free periods after checking the latest rainfall forecast. Monitor crops regularly for pests and diseases, particularly under humid conditions. Keep livestock in clean, dry and well-ventilated shelters and protect feed and fodder from rain and moisture. Conserve rainwater where possible and maintain farm ponds and other water storage structures. Follow weather updates through the Meghdoot app and lightning alerts through the Damini app.

SMS Advisory:

To prevent African swine fever, restrict outsiders' entry into pig farms. Keep the farm, pens and equipment clean and disinfect regularly. Provide clean water and balanced feed.

Crop Specific Advisory:

Crop (Stage)	Crop Specific Advisory
RICE (Millking)	Considering the milky stage of rice, farmers are advised to apply muriate of potash (MOP) and micronutrient mixture through foliar spraying as per recommended doses. Maintain a water level of 2–3 cm in the rice field. Regularly monitor the crop for pest and disease incidence. If Gandhi bug infestation is observed, spray neem seed kernel extract at 5% concentration. Always check the latest weather forecast before applying fertilizers or plant protection chemicals, and avoid spraying during rainfall.
NUTMEG (Flowering)	Considering the emergence of new flushes in clove and nutmeg, farmers are advised to clean the basins of the plants and apply coconut husk mulch to conserve soil moisture and support bud development.

Horticulture Specific Advisory:

Horticulture (Stage)	Horticulture Specific Advisory
BLACK PEPPER (Vegetative)	In black pepper, if yellowing of leaf veins is observed, farmers are advised to apply neem cake @ 500 g per plant and drench the soil with 1% Bordeaux mixture. For preparing 100 litres of 1% Bordeaux mixture, dissolve 1 kg copper sulphate in 50 litres of water and 1 kg fresh lime separately in 50 litres of water; slowly mix the copper sulphate solution into the lime suspension with continuous stirring, filter and use the mixture fresh.
BRINJAL (Fruit formation)	Farmers are advised to regularly inspect brinjal fields. If Bacterial wilt symptoms observed, remove and destroy the infected plants to prevent the spread of the disease. Practise Crop rotaion with non solanaceous crops Grow bacterial wilt-resistant brinjal varieties such as Dweep CARI Brinjal-1. Apply 5% bleaching powder as a soil drench and use streptocycline 2% for seed treatment.
COCONUT	Farmers who have not yet applied post-monsoon fertilizers are advised to apply well-decomposed organic manure along with 50% of the recommended dose of fertilizers (RDF) to coconut, arecanut and other perennial crops. Apply fertilizers around the palm basin and incorporate them into the soil during rain-free periods to minimize nutrient loss. Always check the latest weather forecast before fertilizer application and avoid applying fertilizers when rainfall is expected.
GINGER (Grand growth)	As ginger, turmeric and mango ginger crops are in the rhizome development stage, farmers are advised to carry out earthing up and mulching to cover exposed rhizomes, conserve soil moisture and promote healthy rhizome development. Use organic mulch such as dry leaves or crop residues, and ensure proper drainage to prevent waterlogging.

Live Stock Specific Advisory:

Live Stock	Live Stock Specific Advisory
PIG	Considering the risk of African Swine Fever (ASF) transmission, farmers are advised to restrict the entry of outsiders into pig farms and maintain proper farm hygiene by keeping the surroundings clean and regularly cleaning and disinfecting pig pens and equipment. Provide pigs with good-quality, balanced feed and clean drinking water in adequate quantities, and avoid feeding contaminated feed or water. Healthy pigs should be dewormed at regular intervals as recommended by the local veterinarian and provided with appropriate vitamin and mineral supplements to maintain their health and improve overall productivity.

Live Stock	Live Stock Specific Advisory
COW	Farmers are advised to provide livestock with clean and properly filtered drinking water. Animals should not be allowed to bathe or wallow in stagnant water to prevent the spread of waterborne diseases. Animal dipping should be carried out during daytime under suitable weather conditions. Regular deworming should also be undertaken using appropriate deworming medicines as per veterinary advice.

Likely impacts of weather warnings (General)

There is no weather warning for the upcoming five days. No significant weather-related impact or damage is expected during this period.

Impact based advisories (General)

No impact-based advisory is provided as there is no weather warning.Carry out routine household, workplace and outdoor activities as planned. Keep checking the latest weather forecast and warnings and take necessary precautions if the weather situation changes.

Farmers are advised to download Unified “Mausam” and "Meghdoot" android application on mobile for Weather forecast and weather based Agromet Advisories and "Damini" android application for forecast of Thunderstorm and lightening.

Mausam MobileApp link: https://play.google.com/store/apps/details/

Meghdoot MobileApp link: https://play.google.com/store/apps/details

Damini MobileApp link : https://play.google.com/store/apps/details

ग्रामीण कृषि मौसम सेवा
भारत मौसम विज्ञान विभाग
केंद्रीय द्वीपीय कृषि अनुसंधान संस्थान
पोर्टब्लेयर, अंडमान तथा निकोबार द्वीपसमूह

मौसम आधारित कृषि परामर्श सेवाएं

दिनांक : 03-11-2026

N-AND-M-ANDAMAN(पश्चिम वङ्ग) के मौसम का पूर्वानुमान - जारी करनेका दिन :2026-11-03 (अगले 5 दिनों के 8:30 IST तक वैध)

मौसम कारक	2026-11-04	2026-11-05	2026-11-06	2026-11-07	2026-11-08
वर्षा (मिमी)	3.0	3.0	7.0	8.0	5.0
अधिकतम तापमान(से.)	30.0	30.0	30.0	30.0	30.0
न्यूनतम तापमान(से.)	25.0	25.0	25.0	25.0	25.0
अधिकतम सापेक्षिक आर्द्रता (%)	89	89	88	86	79
न्यूनतम सापेक्षिक आर्द्रता (%)	69	71	68	70	57
हवा की गति (किमी प्रति घंटा)	10	7	3	5	6
पवन दिशा (डिग्री)	9	32	41	13	32
क्लाउड कवर (ओक्टा)	5	3	4	3	6
चेतावनी	कोई चेतावनी नहीं	कोई चेतावनी नहीं	कोई चेतावनी नहीं	कोई चेतावनी नहीं	कोई चेतावनी नहीं

पूर्वानुमान सारांश:

04 नवंबर 2026 से शुरू होने वाले अगले पाँच दिनों के दौरान हल्की वर्षा होने की संभावना है। वर्षा की मात्रा 3.00 से 8.00 मिमी के बीच रहने तथा कुल संचयी वर्षा 26.00 मिमी होने की संभावना है। अधिकतम तापमान लगभग 30.0°C तथा न्यूनतम तापमान लगभग 25.0°C रहने की संभावना है। अधिकतम सापेक्ष आर्द्रता 79% से 89% तथा न्यूनतम सापेक्ष आर्द्रता 57% से 71% के बीच रहने की संभावना है। हवा की गति 3 से 10 किमी प्रति घंटा रहने तथा हवा की दिशा उत्तर (9°) से उत्तर-पूर्व (41°) के बीच परिवर्तित होने की संभावना है। आसमान में आंशिक रूप से बादल छाए रहने से लेकर सामान्यतः बादल छाए रहने की संभावना है।

मौसम चेतावनियाँ (अगले दिन के 08:30 IST तक मान्य):

आगामी पाँच दिनों के लिए कोई मौसम चेतावनी नहीं है।

मौसम चेतावनियों के कृषि पर संभावित प्रभाव और संबंधित एग्रोमेट सलाह:

किसी मौसम चेतावनी के जारी नहोनेके कारण कोई महत्वपूर्ण क्षतिहोने की संभावना नहीं है।

सामान्य सलाहकार:

हल्की वर्षा के मौसम पूर्वानुमान को ध्यान में रखते हुए किसानों को सलाह दी जाती है कि संभावित वर्षा का उपयोग सिंचाई प्रबंधन की योजना बनाने में करें तथा जहां मिट्टी में पर्याप्त नमी उपलब्ध हो, वहां सिंचाई से बचें। उर्वरकों एवं पौध संरक्षण रसायनों का प्रयोग नवीनतम मौसम पूर्वानुमान की जानकारी लेने के बाद उपयुक्त वर्षा रहित अवधि में ही करें। आर्द्र परिस्थितियों में कीट एवं रोगों की संभावना को देखते हुए फसलों की नियमित निगरानी करें। पशुओं को स्वच्छ, सूखे एवं हवादार आश्रय स्थलों में रखें तथा चारा एवं पशु आहार को वर्षा एवं नमी से सुरक्षित रखें। वर्षा जल का अधिकतम संचयन करें तथा खेतों में बने तालाबों एवं अन्य जल भंडारण संरचनाओं का उचित रखरखाव करें। मौसम संबंधी जानकारी के लिए मेघदूत (Meghdoot) ऐप तथा आकाशीय बिजली से संबंधित चेतावनियों के लिए दामिनी (Damini) ऐप का नियमित उपयोग करें।

लघु संदेश सलाहकार:

African Swine Fever से बचाव हेतु सूअर फार्म में बाहरी व्यक्तियों का प्रवेश सीमित रखें। फार्म, बाड़ों एवं उपकरणों की नियमित सफाई एवं कीटाणुशोधन करें। सूअरों को स्वच्छ पानी एवं संतुलित आहार प्रदान करें।

फ़सल विशिष्ट सलाह:

फ़सल	फ़सल विशिष्ट सलाह
चावल	धान की फसल की दूधिया अवस्था को ध्यान में रखते हुए, किसानों को सलाह दी जाती है कि अनुशंसित मात्रा के अनुसार म्यूरेट ऑफ पोटाश (MOP) एवं सूक्ष्म पोषक तत्व मिश्रण का पर्णीय छिड़काव करें। धान के खेत में 2–3 सेमी जलस्तर बनाए रखें। फसल में कीट एवं रोगों के प्रकोप की नियमित निगरानी करें। यदि गांधी बग का प्रकोप दिखाई दे, तो 5% सांद्रता वाले नीम बीज गिरी अर्क का छिड़काव करें। उर्वरकों एवं पौध संरक्षण रसायनों का प्रयोग करने से पहले हमेशा नवीनतम मौसम पूर्वानुमान की जानकारी प्राप्त करें तथा वर्षा के दौरान छिड़काव करने से बचें।
जायफल	लौंग एवं जायफल के पौधों में नई कोंपलों के निकलने को ध्यान में रखते हुए किसानों को सलाह दी जाती है कि पौधों के थालों की सफाई करें तथा मिट्टी में नमी बनाए रखने और कलियों के समुचित विकास को बढ़ावा देने के लिए नारियल की भूसी से मल्चिंग करें।

बागवानी विशिष्ट सलाह:

बागवानी	बागवानी विशिष्ट सलाह
काली मिर्च	काली मिर्च में पत्तियों की शिराओं के पीले पड़ने के लक्षण दिखाई देने पर किसानों को सलाह दी जाती है कि प्रत्येक पौधे में 500 ग्राम नीम की खली डालें तथा 1% बोर्डो मिश्रण से मिट्टी की सिंचाई (ड्रेंचिंग) करें। 100 लीटर 1% बोर्डो मिश्रण तैयार करने के लिए 1 किलोग्राम कॉपर सल्फेट को 50 लीटर पानी में तथा 1 किलोग्राम ताजे बुझा हुए चूने को अलग से 50 लीटर पानी में घोलें। इसके बाद लगातार हिलाते हुए कॉपर सल्फेट के घोल को धीरे-धीरे चूने के घोल में मिलाएं। मिश्रण को छानकर ताजा ही उपयोग करें।
बैंगन	किसानों को सलाह दी जाती है कि बैंगन के खेतों का नियमित निरीक्षण करें। यदि Bacterial wilt के लक्षण दिखाई दें, तो रोग के प्रसार को रोकने के लिए संक्रमित पौधों को उखाड़कर नष्ट कर दें। अन्य कुलों की सब्जियों वाली फसलों के साथ फसल चक्र अपनाएँ तथा जीवाणु म्लानि प्रतिरोधी बैंगन की किस्मों, जैसे Dweep CARI Brinjal-1, की खेती करें। मिट्टी में 5% ब्लीचिंग पाउडर (Bleaching Powder) का घोल डालें तथा बीज उपचार के लिए 2% Streptocyclin का उपयोग करें।
नारियल	जिन किसानों ने अभी तक मानसूनोत्तर उर्वरकों का प्रयोग नहीं किया है, उन्हें सलाह दी जाती है कि वे नारियल, सुपारी एवं अन्य बहुवर्षीय फसलों में अच्छी तरह सड़ी हुई जैविक खाद के साथ अनुशंसित उर्वरक मात्रा (RDF) का 50% प्रयोग करें। उर्वरकों को पौधों के थालों के चारों ओर डालकर मिट्टी में अच्छी तरह मिला दें। पोषक तत्वों के नुकसान को कम करने के लिए उर्वरकों का प्रयोग वर्षा रहित अवधि में ही करें। उर्वरकों के प्रयोग से पूर्व नवीनतम मौसम पूर्वानुमान अवश्य देखें तथा वर्षा की संभावना होने पर उर्वरकों का प्रयोग न करें।
अदरक	अदरक, हल्दी एवं आम-अदरक की फसलें वर्तमान में प्रकंद विकास अवस्था में हैं। अतः किसानों को सलाह दी जाती है कि खुले हुए प्रकंदों को मिट्टी से ढकने, मृदा में नमी बनाए रखने तथा प्रकंदों के समुचित विकास को बढ़ावा देने के लिए मिट्टी चढ़ाने (अर्थिंग-अप) एवं पलवार (मल्चिंग) की प्रक्रिया अपनाएं। इसके लिए सूखी पत्तियों या फसल अवशेषों जैसे जैविक पदार्थों का उपयोग करें तथा खेत में जलभराव रोकने के लिए उचित जल निकासी की व्यवस्था सुनिश्चित करें।

पशुपालन विशिष्ट सलाह:

पशुपालन	पशुपालन विशिष्ट सलाह
सूअर	अफ्रीकन स्वाइन फीवर (ASF) के संक्रमण के जोखिम को देखते हुए, किसानों को सलाह दी जाती है कि वे सूअरों के फार्म में बाहरी व्यक्तियों के प्रवेश को प्रतिबंधित करें तथा फार्म परिसर को स्वच्छ रखें। सूअरशालाओं एवं उपयोग में आने वाले उपकरणों की नियमित रूप से सफाई एवं कीटाणुशोधन करें। सूअरों को पर्याप्त मात्रा में अच्छी गुणवत्ता वाला संतुलित आहार एवं स्वच्छ पेयजल उपलब्ध कराएं तथा दूषित आहार या पानी देने से बचें। स्वस्थ सूअरों को स्थानीय पशु चिकित्सक की सलाह के अनुसार नियमित अंतराल पर कृमिनाशक दवा दें तथा उनके स्वास्थ्य और उत्पादकता को बनाए रखने के लिए उचित मात्रा में विटामिन एवं खनिज पूरक प्रदान करें।
गाय	पशुपालकों को सलाह दी जाती है कि पशुओं को स्वच्छ एवं उचित रूप से छना हुआ पेयजल उपलब्ध कराएं। जलजनित रोगों के प्रसार को रोकने के लिए पशुओं को ठहरे हुए पानी में न नहाने दें और न ही उन्हें उसमें लोटने दें। पशुओं का डिपिंग (परजीवी नियंत्रण हेतु औषधीय घोल में स्नान) दिन के समय अनुकूल मौसम परिस्थितियों में करें। पशुओं को पशु चिकित्सक की सलाह के अनुसार उपयुक्त कृमिनाशक दवाओं से नियमित रूप से कृमिनाशन कराएं।

मौसम चेतावनियों के संभावित प्रभाव (सामान्य):

आगामी पाँच दिनों के लिए कोई मौसम चेतावनी नहीं है। इस अवधिके दौरान मौसम से संबंधित किसी महत्वपूर्ण प्रभाव याक्षतिकी संभावना नहीं है।

प्रभाव आधारित सलाह (सामान्य):

मौसम संबंधी कोई चेतावनी नहीं होने के कारण कोई प्रभाव-आधारित परामर्श नहीं है।घर, कार्यस्थलऔर बाहर के नियमित कार्य योजनानु सार करें।मौसम की स्थिति में बदलाव होने परआवश्यक सावधानियाँ बरतनेके लिए नवीनतम मौसम पूर्वानुमानऔर चेतावनियों की जानकारी लेते रहें।

Farmers are advised to download Unified ‘Mausam’ and "Meghdoot" android application on mobile for Weather forecast and weather based Agromet Advisories and "Damini" android application for forecast of Thunderstorm and lightening.

Mausam MobileApp link: https://play.google.com/store/apps/details/

Meghdoot MobileApp link: https://play.google.com/store/apps/details

Damini MobileApp link : https://play.google.com/store/apps/details

Gramin Krishi Mausam Sewa
Agromet Advisory Bulletin for NICOBAR District
Central Agricultural Research Institute
Port Blair, Andaman & Nicobar Islands

Agromet Advisory Bulletin

Date : 2026-11-03

Weather Forecast of District NICOBAR (West Bengal) Issued On : 2026-11-03 (Valid Till 08:30 IST of the next 5 days)

Parameter	2026-11-04	2026-11-05	2026-11-06	2026-11-07	2026-11-08
Rainfall(mm)	4.0	7.0	10.0	5.0	7.0
Tmax(°C)	31.0	31.0	31.0	31.0	31.0
Tmin(°C)	27.0	27.0	27.0	27.0	27.0
RH-I(%)	91	91	89	89	83
RH-II(%)	75	77	70	73	66
Wind Speed(kmph)	10	11	2	5	4
Wind Direction(Degree)	39	61	100	348	17
Cloud Cover(Octa)	6	7	5	7	7
Warning	No Warning	No Warning	No Warning	No Warning	No Warning

Forecast Summary:

The weather forecast for the next five days, starting from 04 November 2026, indicates that light rainfall is expected, with rainfall amounts ranging from 4.00 to 10.00 mm and cumulative rainfall expected to be 33.00 mm. The maximum temperature is expected to be around 31.0°C, while the minimum temperature will be around 27.0°C. Maximum relative humidity is likely to range from 83% to 91%, and minimum relative humidity from 66% to 77%. Wind speed is expected to vary between 2 and 11 kmph, with wind directions varying from north-northeast (17°) to north-northwest (348°). Cloud conditions are expected to remain generally cloudy.

Weather Warnings (Valid Till 08:30 IST of the next day)

There is no weather warning for upcoming five days.

Likely impacts of weather warnings on Agriculture and associated Agromet advisories

Since no warnings have been issued, no significant damage is expected.

General Advisory:

Considering the forecast of light rainfall, farmers are advised to use the expected rainfall for planning irrigation and avoid irrigation where adequate soil moisture is available. Fertilizer and plant protection applications may be taken up during suitable rain-free periods after checking the latest rainfall forecast. Monitor crops regularly for pests and diseases, particularly under humid conditions. Keep livestock in clean, dry and well-ventilated shelters and protect feed and fodder from rain and moisture. Conserve rainwater where possible and maintain farm ponds and other water storage structures. Follow weather updates through the Meghdoot app and lightning alerts through the Damini app.

SMS Advisory:

To prevent African swine fever, restrict outsiders' entry into pig farms. Keep the farm, pens and equipment clean and disinfect regularly. Provide clean water and balanced feed.

Crop Specific Advisory:

Crop (Stage)	Crop Specific Advisory
NUTMEG (Flowering)	Considering the emergence of new flushes in clove and nutmeg, farmers are advised to clean the basins of the plants and apply coconut husk mulch to conserve soil moisture and support bud development.

Horticulture Specific Advisory:

Horticulture (Stage)	Horticulture Specific Advisory
BLACK PEPPER (Vegetative)	In black pepper, if yellowing of leaf veins is observed, farmers are advised to apply neem cake @ 500 g per plant and drench the soil with 1% Bordeaux mixture. For preparing 100 litres of 1% Bordeaux mixture, dissolve 1 kg copper sulphate in 50 litres of water and 1 kg fresh lime separately in 50 litres of water; slowly mix the copper sulphate solution into the lime suspension with continuous stirring, filter and use the mixture fresh.
BRINJAL (Fruit formation)	Farmers are advised to regularly inspect brinjal fields. If Bacterial wilt symptoms observed, remove and destroy the infected plants to prevent the spread of the disease. Practise Crop rotaion with non solanaceous crops Grow bacterial wilt-resistant brinjal varieties such as Dweep CARI Brinjal-1. Apply 5% bleaching powder as a soil drench and use streptocycline 2% for seed treatment.
COCONUT	Farmers who have not yet applied post-monsoon fertilizers are advised to apply well-decomposed organic manure along with 50% of the recommended dose of fertilizers (RDF) to coconut, arecanut and other perennial crops. Apply fertilizers around the palm basin and incorporate them into the soil during rain-free periods to minimize nutrient loss. Always check the latest weather forecast before fertilizer application and avoid applying fertilizers when rainfall is expected.
GINGER (Grand growth)	As ginger, turmeric and mango ginger crops are in the rhizome development stage, farmers are advised to carry out earthing up and mulching to cover exposed rhizomes, conserve soil moisture and promote healthy rhizome development. Use organic mulch such as dry leaves or crop residues, and ensure proper drainage to prevent waterlogging.

Live Stock Specific Advisory:

Live Stock	Live Stock Specific Advisory
PIG	Considering the risk of African Swine Fever (ASF) transmission, farmers are advised to restrict the entry of outsiders into pig farms and maintain proper farm hygiene by keeping the surroundings clean and regularly cleaning and disinfecting pig pens and equipment. Provide pigs with good-quality, balanced feed and clean drinking water in adequate quantities, and avoid feeding contaminated feed or water. Healthy pigs should be dewormed at regular intervals as recommended by the local veterinarian and provided with appropriate vitamin and mineral supplements to maintain their health and improve overall productivity.
GOAT	Farmers are advised to provide livestock with clean and properly filtered drinking water. Animals should not be allowed to bathe or wallow in stagnant water to prevent the spread of waterborne diseases. Animal dipping should be carried out during daytime under suitable weather conditions. Regular deworming should also be undertaken using appropriate deworming medicines as per veterinary advice.

Likely impacts of weather warnings (General)

There is no weather warning for the upcoming five days. No significant weather-related impact or damage is expected during this period.

Impact based advisories (General)

No impact-based advisory is provided as there is no weather warning.Carry out routine household, workplace and outdoor activities as planned. Keep checking the latest weather forecast and warnings and take necessary precautions if the weather situation changes.

Farmers are advised to download Unified “Mausam” and "Meghdoot" android application on mobile for Weather forecast and weather based Agromet Advisories and "Damini" android application for forecast of Thunderstorm and lightening.

Mausam MobileApp link: https://play.google.com/store/apps/details/

Meghdoot MobileApp link: https://play.google.com/store/apps/details

Damini MobileApp link : https://play.google.com/store/apps/details

ग्रामीण कृषि मौसम सेवा
भारत मौसम विज्ञान विभाग
केंद्रीय द्वीपीय कृषि अनुसंधान संसथान
पोर्टब्लेयर, अंडमान तथा निकोबार द्वीपसमूह

मौसम आधारित कृषि परामर्श सेवाएं

दिनांक : 03-11-2026

NICOBAR(पश्चिम बङ्ग) के मौसम का पूर्वानुमान - जारी करनेका दिन :2026-11-03 (अगले 5 दिनों के 8:30 IST तक वैध)

मौसम कारक	2026-11-04	2026-11-05	2026-11-06	2026-11-07	2026-11-08
वर्षा (मिमी)	4.0	7.0	10.0	5.0	7.0
अधिकतम तापमान(से.)	31.0	31.0	31.0	31.0	31.0
न्यूनतम तापमान(से.)	27.0	27.0	27.0	27.0	27.0
अधिकतम सापेक्षिक आर्द्रता (%)	91	91	89	89	83
न्यूनतम सापेक्षिक आर्द्रता (%)	75	77	70	73	66
हवा की गति (किमी प्रति घंटा)	10	11	2	5	4
पवन दिशा (डिग्री)	39	61	100	348	17
क्लाउड कवर (ओक्टा)	6	7	5	7	7
चेतावनी	कोई चेतावनी नहीं	कोई चेतावनी नहीं	कोई चेतावनी नहीं	कोई चेतावनी नहीं	कोई चेतावनी नहीं

पूर्वानुमान सारांश:

04 नवंबर 2026 से शुरू होने वाले अगले पाँच दिनों के दौरान हल्की वर्षा होने की संभावना है। वर्षा की मात्रा 4.00 से 10.00 मिमी के बीच रहने तथा कुल संचयी वर्षा 33.00 मिमी होने की संभावना है। अधिकतम तापमान लगभग 31.0°C तथा न्यूनतम तापमान लगभग 27.0°C रहने की संभावना है। अधिकतम सापेक्ष आर्द्रता 83% से 91% तथा न्यूनतम सापेक्ष आर्द्रता 66% से 77% के बीच रहने की संभावना है। हवा की गति 2 से 11 किमी प्रति घंटा रहने तथा हवा की दिशा उत्तर-उत्तर-पूर्व (17°) से उत्तर-उत्तर-पश्चिम (348°) के बीच परिवर्तित होने की संभावना है। आसमान में सामान्यतः बादल छाए रहने की संभावना है।

मौसम चेतावनियाँ (अगले दिन के 08:30 IST तक मान्य):

आगामी पाँच दिनों के लिए कोई मौसम चेतावनी नहीं है।

मौसम चेतावनियों के कृषि पर संभावित प्रभाव और संबंधित एग्रोमेट सलाह:

किसी मौसम चेतावनी के जारी नहोनेके कारण कोई महत्वपूर्ण क्षतिहोने की संभावना नहीं है।

सामान्य सलाहकार:

हल्की वर्षा के मौसम पूर्वानुमान को ध्यान में रखते हुए किसानों को सलाह दी जाती है कि संभावित वर्षा का उपयोग सिंचाई प्रबंधन की योजना बनाने में करें तथा जहां मिट्टी में पर्याप्त नमी उपलब्ध हो, वहां सिंचाई से बचें। उर्वरकों एवं पौध संरक्षण रसायनों का प्रयोग नवीनतम मौसम पूर्वानुमान की जानकारी लेने के बाद उपयुक्त वर्षा रहित अवधि में

ही करें। आर्द्र परिस्थितियों में कीट एवं रोगों की संभावना को देखते हुए फसलों की नियमित निगरानी करें। पशुओं को स्वच्छ, सूखे एवं हवादार आश्रय स्थलों में रखें तथा चारा एवं पशु आहार को वर्षा एवं नमी से सुरक्षित रखें। वर्षा जल का अधिकतम संचयन करें तथा खेतों में बने तालाबों एवं अन्य जल भंडारण संरचनाओं का उचित रखरखाव करें। मौसम संबंधी जानकारी के लिए मेघदूत (Meghdoot) ऐप तथा आकाशीय बिजली से संबंधित चेतावनियों के लिए दामिनी (Damini) ऐप का नियमित उपयोग करें।

लघु संदेश सलाहकार:

African Swine Fever से बचाव हेतु सूअर फार्म में बाहरी व्यक्तियों का प्रवेश सीमित रखें। फार्म, बाड़ों एवं उपकरणों की नियमित सफाई एवं कीटाणुशोधन करें। सूअरों को स्वच्छ पानी एवं संतुलित आहार प्रदान करें।

फ़सल विशिष्ट सलाह:

फ़सल	फ़सल विशिष्ट सलाह
जायफल	लौंग एवं जायफल के पौधों में नई कोंपलों के निकलने को ध्यान में रखते हुए किसानों को सलाह दी जाती है कि पौधों के थालों की सफाई करें तथा मिट्टी में नमी बनाए रखने और कलियों के समुचित विकास को बढ़ावा देने के लिए नारियल की भूसी से मल्चिंग करें।

बागवानी विशिष्ट सलाह:

बागवानी	बागवानी विशिष्ट सलाह
काली मिर्च	काली मिर्च में पत्तियों की शिराओं के पीले पड़ने के लक्षण दिखाई देने पर किसानों को सलाह दी जाती है कि प्रत्येक पौधे में 500 ग्राम नीम की खली डालें तथा 1% बोर्डो मिश्रण से मिट्टी की सिंचाई (ड्रेंचिंग) करें। 100 लीटर 1% बोर्डो मिश्रण तैयार करने के लिए 1 किलोग्राम कॉपर सल्फेट को 50 लीटर पानी में तथा 1 किलोग्राम ताजे बुझा हुए चूने को अलग से 50 लीटर पानी में घोलें। इसके बाद लगातार हिलाते हुए कॉपर सल्फेट के घोल को धीरे-धीरे चूने के घोल में मिलाएं। मिश्रण को छानकर ताजा ही उपयोग करें।
बैंगन	किसानों को सलाह दी जाती है कि बैंगन के खेतों का नियमित निरीक्षण करें। यदि Bacterial wilt के लक्षण दिखाई दें, तो रोग के प्रसार को रोकने के लिए संक्रमित पौधों को उखाड़कर नष्ट कर दें। अन्य कुलों की सब्जियों वाली फसलों के साथ फसल चक्र अपनाएँ तथा जीवाणु म्लानि प्रतिरोधी बैंगन की किस्मों, जैसे Dweep CARI Brinjal-1, की खेती करें। मिट्टी में 5% ब्लीचिंग पाउडर (Bleaching Powder) का घोल डालें तथा बीज उपचार के लिए 2% Streptocyclin का उपयोग करें।
नारियल	जिन किसानों ने अभी तक मानसूनोत्तर उर्वरकों का प्रयोग नहीं किया है, उन्हें सलाह दी जाती है कि वे नारियल, सुपारी एवं अन्य बहुवर्षीय फसलों में अच्छी तरह सड़ी हुई जैविक खाद के साथ अनुशंसित उर्वरक मात्रा (RDF) का 50% प्रयोग करें। उर्वरकों को पौधों के थालों के चारों ओर डालकर मिट्टी में अच्छी तरह मिला दें। पोषक तत्वों के नुकसान को कम करने के लिए उर्वरकों का प्रयोग वर्षा रहित अवधि में ही करें। उर्वरकों के प्रयोग से पूर्व नवीनतम मौसम पूर्वानुमान अवश्य देखें तथा वर्षा की संभावना होने पर उर्वरकों का प्रयोग न करें।
अदरक	अदरक, हल्दी एवं आम-अदरक की फसलें वर्तमान में प्रकंद विकास अवस्था में हैं। अतः किसानों को सलाह दी जाती है कि खुले हुए प्रकंदों को मिट्टी से ढकने, मृदा में नमी बनाए रखने तथा प्रकंदों के समुचित विकास को बढ़ावा देने के लिए मिट्टी चढ़ाने (अर्थिंग-अप) एवं पलवार (मल्चिंग) की प्रक्रिया अपनाएं। इसके लिए सूखी पत्तियों या फसल अवशेषों जैसे जैविक पदार्थों का उपयोग करें तथा खेत में जलभराव रोकने के लिए उचित जल निकासी की व्यवस्था सुनिश्चित करें।

पशुपालन विशिष्ट सलाह:

पशुपालन	पशुपालन विशिष्ट सलाह
सूअर	अफ्रीकन स्वाइन फीवर (ASF) के संक्रमण के जोखिम को देखते हुए, किसानों को सलाह दी जाती है कि वे सूअरों के फार्म में बाहरी व्यक्तियों के प्रवेश को प्रतिबंधित करें तथा फार्म परिसर को स्वच्छ रखें। सूअरशालाओं एवं उपयोग में आने वाले उपकरणों की नियमित रूप से सफाई एवं कीटाणुशोधन करें। सूअरों को पर्याप्त मात्रा में अच्छी गुणवत्ता वाला संतुलित आहार एवं स्वच्छ पेयजल उपलब्ध कराएं तथा दूषित आहार या पानी देने से बचें। स्वस्थ सूअरों को स्थानीय पशु चिकित्सक की सलाह के अनुसार नियमित अंतराल पर कृमिनाशक दवा दें तथा उनके स्वास्थ्य और उत्पादकता को बनाए रखने के लिए उचित मात्रा में विटामिन एवं खनिज पूरक प्रदान करें।

पशुपालन	पशुपालन विशिष्ट सलाह
बकरा	पशुपालकों को सलाह दी जाती है कि पशुओं को स्वच्छ एवं उचित रूप से छना हुआ पेयजल उपलब्ध कराएं। जलजनित रोगों के प्रसार को रोकने के लिए पशुओं को ठहरे हुए पानी में न नहाने दें और न ही उन्हें उसमें लोटने दें। पशुओं का डिपिंग (परजीवी नियंत्रण हेतु औषधीय घोल में स्नान) दिन के समय अनुकूल मौसम परिस्थितियों में करें। पशुओं को पशु चिकित्सक की सलाह के अनुसार उपयुक्त कृमिनाशक दवाओं से नियमित रूप से कृमिनाशन कराएं।

मौसम चेतावनियों के संभावित प्रभाव (सामान्य):

आगामी पाँच दिनों के लिए कोई मौसम चेतावनी नहीं है। इस अवधिके दौरान मौसम से संबंधित किसी महत्वपूर्ण प्रभाव याक्षतिकी संभावना नहीं है।

प्रभाव आधारित सलाह (सामान्य):

मौसम संबंधी कोई चेतावनी नहीं होने के कारण कोई प्रभाव-आधारित परामर्श नहीं है।घर, कार्यस्थलऔर बाहर के नियमित कार्य योजनानु सार करें।मौसम की स्थिति में बदलाव होने परआवश्यक सावधानियाँ बरतनेके लिए नवीनतम मौसम पूर्वानुमानऔर चेतावनियों की जानकारी लेते रहें।

Farmers are advised to download Unified ﷺ and "Meghdoot" android application on mobile for Weather forecast and weather based Agromet Advisories and "Damini" android application for forecast of Thunderstorm and lightening.

Mausam MobileApp link: https://play.google.com/store/apps/details/

Meghdoot MobileApp link: https://play.google.com/store/apps/details

Damini MobileApp link : https://play.google.com/store/apps/details